

Hyperplane sections and stable derived categories

Kazushi Ueda

Abstract

We discuss the relation between the graded stable derived category of a hypersurface and that of its hyperplane section. The motivation comes from the compatibility between homological mirror symmetry for the Calabi-Yau manifold defined by an invertible polynomial and that for the singularity defined by the same polynomial.

1 Introduction

Let $(Y, \mathcal{O}_Y(1))$ be a polarized smooth projective variety of dimension d over a field $\mathbf{k}$, and $X = s^{-1}(0)$ be a smooth hypersurface defined by a section $s \in H^0(\mathcal{O}_Y(a))$ of degree a . For coherent sheaves $\mathcal{E}$ and $\mathcal{E}'$ on X restricted from those on Y , the push-forward functor $\iota_* : D^b \text{coh } X \rightarrow D^b \text{coh } Y$ along the inclusion $\iota : X \hookrightarrow Y$ satisfies

$$\text{Hom}_Y^i(\iota_* \mathcal{E}, \iota_* \mathcal{E}') \cong \text{Hom}_X^i(\mathcal{E}, \mathcal{E}') \oplus \text{Hom}_X^{d-i}(\mathcal{E}', \mathcal{E} \otimes \omega_Y|_X)^\vee, \quad (1.1)$$

where $\omega_Y|_X = \iota^* \omega_Y$ is the restriction of the dualizing sheaf of Y to X and $\bullet^\vee$ denotes the $\mathbf{k}$ -dual vector space. In particular, if Y has the trivial dualizing sheaf, then one has

$$\text{Hom}_Y^i(\iota_* \mathcal{E}, \iota_* \mathcal{E}') \cong \text{Hom}_X^i(\mathcal{E}, \mathcal{E}') \oplus \text{Hom}_X^{d-i}(\mathcal{E}', \mathcal{E}).$$

If $\omega_Y \cong \mathcal{O}_Y(r)$ for some $r \in \mathbb{Z}$ and $H^i(\mathcal{O}_Y(n)) = 0$ for any $n \in \mathbb{Z}$ unless $i = 0, d$, then the graded ring $\overline{S} = \bigoplus_{n=0}^\infty H^0(\mathcal{O}_Y(n))$ is *Gorenstein* with *a-invariant* r , in the sense that S has finite injective dimension as a graded S -module and the graded canonical module K_S is isomorphic to the free module $S(r)$ [Orl09, Lemma 2.12]. The latter condition is equivalent to the isomorphism $\mathbb{R}\text{Hom}_{\overline{S}}(\mathbf{k}, \overline{S}) \cong \mathbf{k}(-r)[-d-1]$. Here the round bracket and the square bracket indicate the shift in the *internal* and *homological* grading respectively.

The *graded stable derived category* of $\overline{S}$ is defined as the quotient category

$$D_{\text{sing}}^b(\text{gr } \overline{S}) := D^b(\text{gr } \overline{S}) / D^{\text{perf}}(\text{gr } \overline{S})$$

of the bounded derived category of finitely-generated graded $\overline{S}$ -modules by the full triangulated subcategory $D^{\text{perf}}(\text{gr } \overline{S})$ consisting of perfect complexes (i.e., bounded complexes of finitely-generated projective modules) [Buc87, Hap91, Kra05, Orl04]. Orlov [Orl09, Theorem 2.13] has shown the existence of

- a full and faithful functor $\Psi_{\overline{S}} : D_{\text{sing}}^b(\text{gr } \overline{S}) \hookrightarrow D^b \text{coh } Y$ if $r < 0$,
- an equivalence $\Psi_{\overline{S}} : D_{\text{sing}}^b(\text{gr } \overline{S}) \xrightarrow{\sim} D^b \text{coh } Y$ if $r = 0$, and

- a full and faithful functor $\Psi_{\overline{S}} : D^b \text{coh } Y \hookrightarrow D_{\text{sing}}^b(\text{gr } \overline{S})$ if $r > 0$.

The graded ring $\overline{R} = \bigoplus_{n=0}^{\infty} H^0(\mathcal{O}_X(n))$ is the quotient ring of $\overline{S}$ by the principal ideal generated by s . Let $\Phi_{\text{gr}} : \text{gr } \overline{R} \rightarrow \text{gr } \overline{S}$ be the functor sending a graded $\overline{R}$ -module to the same module considered as a graded $\overline{S}$ -module. Since $\overline{R}$ is perfect as an $\overline{S}$ -module, the functor Φ_{gr} induces the push-forward functor

$$\Phi_{\text{sing}} : D_{\text{sing}}^b(\text{gr } \overline{R}) \rightarrow D_{\text{sing}}^b(\text{gr } \overline{S})$$

in the stable derived categories. When the a -invariant of $\overline{R}$ is 1 and Y is Calabi-Yau, then [KMU12, Theorem 1.1] states that the functor

$$\Psi_{\overline{S}} \circ \Phi_{\text{sing}} \circ \Psi_{\overline{R}} : D^b \text{coh } X \rightarrow D^b \text{coh } Y$$

is isomorphic to the push-forward functor

$$\iota_* : D^b \text{coh } X \rightarrow D^b \text{coh } Y.$$

In particular, if two objects $\mathcal{E}, \mathcal{E}'$ in $D_{\text{sing}}^b(\text{gr } \overline{R})$ are in the image of $\Psi_{\overline{R}}$, then (1.1) implies

$$\text{Hom}^i(\Phi_{\text{sing}}(\mathcal{E}), \Phi_{\text{sing}}(\mathcal{E}')) \cong \text{Hom}^i(\mathcal{E}, \mathcal{E}') \oplus \text{Hom}^{d-i}(\mathcal{E}', \mathcal{E})^\vee. \quad (1.2)$$

On the other hand, the semiorthogonal complement of the image $\Psi_{\overline{R}}$ is generated by the structure sheaf $\overline{R}/\mathfrak{m}_{\overline{R}}$ of the origin, which goes to $\mathcal{O}_Y[d]$ by the composition $\Psi_{\overline{S}} \circ \Phi_{\text{sing}}$. Since $\overline{R}/\mathfrak{m}_{\overline{R}}$ is exceptional and $\mathcal{O}_Y[d]$ is spherical, (1.2) holds also in this case.

In this paper, we prove a graded stable derived category analog of (1.1) in the following situation: Let R be a graded regular ring with a -invariant r and Krull dimension d , and $S = R \otimes_{\mathbf{k}} \mathbf{k}[w]$ be the tensor product of R with the polynomial ring in one variable w of degree a . We identify the ring R with its image by the natural injection $R \hookrightarrow S$. Let further $f \in R_h$ be a homogeneous element of R of degree h , and $F = f + wg \in S_h$ be a homogeneous element of S of the same degree as f . We will always assume that $g \in wS$. The corresponding quotient rings will be denoted by $\overline{S} = S/(F)$ and $\overline{R} = R/(f) \cong \overline{S}/(w)$.

Theorem 1.1. *For any objects $\mathcal{E}$ and $\mathcal{E}'$ of $D_{\text{sing}}^b(\text{gr } \overline{R})$, one has*

$$\text{Hom}^i(\Phi_{\text{sing}}(\mathcal{E}), \Phi_{\text{sing}}(\mathcal{E}')) \cong \text{Hom}^i(\mathcal{E}, \mathcal{E}') \oplus \text{Hom}^{d-i}(\mathcal{E}', \mathcal{E}(r+h-a))^\vee.$$

In particular, when the a -invariant $r+h$ of $\overline{R}$ and the degree a of the variable w coincides (i.e., when $\overline{S}$ has the trivial canonical module), then one has

$$\text{Hom}^i(\Phi_{\text{sing}}(\mathcal{E}), \Phi_{\text{sing}}(\mathcal{E}')) \cong \text{Hom}^i(\mathcal{E}, \mathcal{E}') \oplus \text{Hom}^{d-i}(\mathcal{E}', \mathcal{E})^\vee. \quad (1.3)$$

The motivation for Theorem 1.1 comes from the compatibility between homological mirror symmetry for the singularity defined by an invertible polynomial and that for the Calabi-Yau manifold defined by the same polynomial. An integer $(d+1) \times (d+1)$ -matrix $(a_{ij})_{i,j=1}^{d+1}$ with non-zero determinant defines a polynomial $f \in \mathbf{k}[x_1, \dots, x_{d+1}]$ by

$$f = \sum_{i=1}^{d+1} x_1^{a_{i1}} \cdots x_{d+1}^{a_{i,d+1}},$$

which is called *invertible* if it has an isolated critical point at the origin. Invertible polynomials play essential role in transposition mirror symmetry of Berglund and Hübsch [BH93], which has attracted much attention recently (see e.g. [Bor, CR11, Kra] and references therein). The quotient ring $\overline{R} = \mathbf{k}[x_1, \dots, x_{d+1}]/(f)$ is naturally graded by the abelian group L generated by $d+2$ elements $\vec{x}_i$ and $\vec{c}$ with relations

$$a_{i1}\vec{x}_1 + \dots + a_{i,d+1}\vec{x}_{d+1} = \vec{c}, \quad i = 1, \dots, d+1.$$

Homological mirror symmetry [Kon95] for invertible polynomials [Tak10] is a conjectural equivalence

$$D_{\text{sing}}^b(\text{gr } \overline{R}) \cong D^b \mathfrak{Fuk} \check{f} \quad (1.4)$$

of triangulated categories. Here $\mathfrak{Fuk} \check{f}$ is the *Fukaya-Seidel category* [Sei08] of the exact symplectic Lefschetz fibration obtained by Morsifying the Berglund-Hübsch transpose $\check{f} = \sum_{i=1}^{d+1} x_1^{a_{1i}} \dots x_{d+1}^{a_{d+1,i}}$ of f . The equivalence (1.4) is proved when $\check{f}$ is the Sebastiani-Thom sum of polynomials of types A or D [FU11, FU].

Assume that one can add one more term to f and obtain another invertible polynomial

$$F = f + x_1^{a_{d+2,1}} \dots x_{d+2}^{a_{d+2,d+2}} \in \mathbf{k}[x_1, \dots, x_{d+2}],$$

with a suitable L -grading on x_{d+2} such that F is homogeneous of degree $\vec{c}$ and the quotient ring $\overline{S} = \mathbf{k}[x_1, \dots, x_{d+2}]/(F)$ is Gorenstein with the trivial a -invariant; $K_{\overline{S}} \cong \overline{S}$.

The zero locus $F^{-1}(0) = \text{Spec } \overline{S}$ has an action of $K = \text{Hom}(L, \mathbb{G}_m)$ coming from the L -grading on $\overline{S}$, and the quotient stack $Y = [(F^{-1}(0) \setminus \mathbf{0})/K]$ of the complement of the origin $\mathbf{0} \in F^{-1}(0)$ by this action is a Calabi-Yau orbifold. The *Berglund-Hübsch transpose* of Y is another Calabi-Yau hypersurface $\check{Y}$ in a weighted projective space defined by the Berglund-Hübsch transpose $\check{F}$ of F . It is conjectured [BH93] that $(Y, \check{Y})$ is a mirror pair, so that there is an equivalence

$$D^b \text{coh } Y \cong D^\pi \mathfrak{Fuk} \check{Y} \quad (1.5)$$

of triangulated categories between the derived category of coherent sheaves on Y and the split-closed derived Fukaya category of $\check{Y}$ [Kon95, Sei11].

The weighted projective hypersurface $\check{Y}$ is a compactification of the Milnor fiber $\check{Y}^\circ = \check{Y} \setminus \{x_{d+2} = 0\} \cong \check{f}^{-1}(-1)$ of $\check{f}$. Let $(V_i)_{i=1}^m$ be a distinguished basis of vanishing cycles of $\check{f}$, considered as Lagrangian submanifolds of $\check{Y}^\circ$. Let further $\mathcal{F}^\rightarrow$ be the directed subcategory of $\mathfrak{Fuk} \check{Y}^\circ$ consisting of $(V_i)_{i=1}^m$; it is an A_∞ -subcategory with

$$\text{hom}_{\mathcal{F}^\rightarrow}(V_i, V_j) = \begin{cases} \mathbf{k} \cdot \text{id}_{V_i} & i = j, \\ \text{hom}_{\mathfrak{Fuk} \check{Y}^\circ}(V_i, V_j) & i < j, \\ 0 & \text{otherwise,} \end{cases}$$

and A_∞ -operations on $\mathcal{F}^\rightarrow$ are inherited from those of $\mathfrak{Fuk} \check{Y}^\circ$ in the obvious way. The Picard-Lefschetz theory [Sei08, Theorem 18.24] gives a derived equivalence

$$D^b \mathcal{F}^\rightarrow \cong D^b \mathfrak{Fuk} \check{f}$$

of $\mathcal{F}^\rightarrow$ and the Fukaya-Seidel category of $\check{f}$. Although the Fukaya category $\mathfrak{Fuk} \check{Y}$ is a deformation of $\mathfrak{Fuk} \check{Y}^\circ$ [Sei02] and the A_∞ -operations on $\mathfrak{Fuk} \check{Y}$ is difficult to compute explicitly, Poincaré duality tells us that the space of morphisms in $\mathfrak{Fuk} \check{Y}$ is given by the *Calabi-Yau completion*

$$\mathrm{Hom}_{\mathfrak{Fuk} \check{Y}}^i(V_j, V_k) = \mathrm{Hom}_{\mathcal{F}^\rightarrow}^i(V_j, V_k) \oplus \mathrm{Hom}_{\mathcal{F}^\rightarrow}^{d-i}(V_k, V_j)^\vee \quad (1.6)$$

of the space of morphisms in $\mathcal{F}^\rightarrow$.

If we assume homological mirror symmetry (1.4) for singularities, then $D_{\mathrm{sing}}^b(\mathrm{gr} \overline{R})$ has a full exceptional collection $(E_i)_{i=1}^m$ corresponding to V_i , and Theorem 1.1 shows that

- the full subcategory of $D_{\mathrm{sing}}^b(\mathrm{gr} \overline{S}) \cong D^b \mathrm{coh} Y$ consisting of $(\Phi_{\mathrm{sing}}(E_i))_{i=1}^m$ is a Calabi-Yau completion of the full subcategory $\mathcal{E}^\rightarrow$ of $D_{\mathrm{sing}}^b(\mathrm{gr} \overline{R})$ consisting of $(E_i)_{i=1}^m$, and
- $\mathcal{E}^\rightarrow$ is the directed subcategory of $D_{\mathrm{sing}}^b(\mathrm{gr} \overline{S})$ consisting of $(\Phi_{\mathrm{sing}}(E_i))_{i=1}^m$.

Now the compatibility of homological mirror symmetry for Calabi-Yau manifolds and that for singularities is the existence of a commutative diagram

$$\begin{array}{ccc} \mathcal{F}^\rightarrow & \hookrightarrow & D^\pi \mathfrak{Fuk} \check{Y} \\ \wr \parallel & & \wr \parallel \\ \mathcal{E}^\rightarrow & \hookrightarrow & D^b \mathrm{coh} Y \end{array} \quad (1.7)$$

where horizontal arrows are embeddings of directed subcategories and vertical equivalences are homological mirror symmetry. Moreover, we expect that the images of the horizontal arrows split-generate the categories on the right, so that one can divide the proof of homological mirror symmetry for Calabi-Yau manifolds into two steps; the first step is the proof of homological mirror symmetry for singularities, and the second step is the analysis of Calabi-Yau completion. This is analogous to the proof of homological mirror symmetry for Calabi-Yau hypersurfaces in projective spaces, which can be interpreted as first proving homological mirror symmetry for the ambient projective space, and then passing to the Calabi-Yau hypersurface by taking a non-trivial Calabi-Yau completion [Sei11, She11, NU].

This paper is organized as follows: In Section 2, we recall basic definitions on matrix factorizations. In Section 3, we give an explicit description of the push-forward in terms of matrix factorizations, which will be used in Section 4 to prove Theorem 1.1.

Acknowledgment: We thank Yukinobu Toda for pointing out an error in (1.1) in the earlier version. This research is supported by Grant-in-Aid for Young Scientists (No. 24740043).

2 Graded matrix factorizations

Let R be a graded regular ring with a -invariant r and Krull dimension d , and $f \in R_h$ be a homogeneous element of degree h . We assume that R is graded by $\mathbb{Z}$ for simplicity of exposition, although our discussion easily generalizes to grading by any abelian group.

A *graded matrix factorization*

$$\begin{array}{ccc} \bigoplus_{i=1}^m R(e_i) & \begin{array}{c} \xrightarrow{\phi} \\ \xleftarrow{\psi} \end{array} & \bigoplus_{i=1}^m R(d_i) \end{array}$$

of f over R consists of

- a pair $M_0 = \bigoplus_{i=1}^m R(d_i)$ and $M_1 = \bigoplus_{i=1}^m R(e_i)$ of graded free R -module and
- a pair (ϕ, ψ) of morphisms $\phi : M_1 \rightarrow M_0$ and $\psi : M_0(-h) \rightarrow M_1$ of graded modules

satisfying $\phi \circ \psi = f \cdot \text{id}_{M_0}$ and $\psi \circ \phi = f \cdot \text{id}_{M_1}$. The integers d_i and e_i are shifts in the internal grading of R -modules, and the natural number $m = \text{rank } M^0 = \text{rank } M^1$ is called the *rank* of the matrix factorization.

A *morphism* of graded matrix factorizations from $(\phi, \psi) : M_1 \rightleftarrows M_0$ to $(\phi', \psi') : M'_1 \rightleftarrows M'_0$ is a pair (α, β) of morphisms $\alpha : M_0 \rightarrow M'_0$ and $\beta : M_1 \rightarrow M'_1$ making the diagram

$$\begin{array}{ccccc} M_0(-h) & \xrightarrow{\psi} & M_1 & \xrightarrow{\phi} & M_0 \\ \alpha \downarrow & & \beta \downarrow & & \alpha \downarrow \\ M'_0(-h) & \xrightarrow{\psi'} & M'_1 & \xrightarrow{\phi'} & M'_0 \end{array}$$

commute; $\alpha \circ \phi = \phi' \circ \beta$ and $\beta \circ \psi = \psi' \circ \alpha$. Here, the morphism $M_0(-h) \rightarrow M'_0(-h)$ of graded R -modules corresponding to the morphism $\alpha : M_0 \rightarrow M'_0$ is denoted by the same symbol.

Two morphisms (α, β) and (α', β') are *homotopic* if there exist morphisms $\xi : M_0 \rightarrow M'_1$ and $\eta : M_1 \rightarrow M'_0$ satisfying $\alpha - \alpha' = \phi' \circ \xi + \eta \circ \psi$ and $\beta - \beta' = \psi' \circ \eta + \xi \circ \phi$.

The *homotopy category of graded matrix factorizations* of f over R is the category $\text{Hmf}_R^{\text{gr}} f$ whose objects are graded matrix factorizations of f over R and whose morphisms are morphisms of graded matrix factorizations up to homotopy.

The category $\text{Hmf}_R^{\text{gr}} f$ is equivalent to the *stable category* $\underline{\text{CM}}^{\text{gr}} \overline{R}$ of graded Cohen-Macaulay modules over the quotient ring $\overline{R} = R/(f)$ [Eis80], which in turn is equivalent to the graded stable derived category $D_{\text{sing}}^b(\text{gr } \overline{R})$ of $\overline{R}$ [Buc87, Orl04].

3 The push-forward of a matrix factorization

We keep the notations from Section 2. Let $S = R \otimes \mathbf{k}[w]$ be another graded ring with $\deg w = a$, and $F = f + wg$ be a homogeneous element of S such that $g \in wS$. The corresponding quotient ring will be denoted by $\overline{S} = S/(F)$, and one has natural injections $R \hookrightarrow S$ and $\overline{R} \hookrightarrow \overline{S}$. The natural ring homomorphism $\overline{S} \rightarrow \overline{S}/(w) \cong \overline{R}$ induces the push-forward functor

$$\Phi_{\text{sing}} : D_{\text{sing}}^b(\text{gr } \overline{R}) \rightarrow D_{\text{sing}}^b(\text{gr } \overline{S}),$$

since $\overline{R}$ is perfect as an $\overline{S}$ -module.

Proposition 3.1. *The functor*

$$\Phi_{\text{Hmf}} : \text{Hmf}_R^{\text{gr}} f \rightarrow \text{Hmf}_S^{\text{gr}} F$$

corresponding to the push-forward functor

$$\Phi_{\text{sing}} : D_{\text{sing}}^b(\text{gr } \overline{R}) \rightarrow D_{\text{sing}}^b(\text{gr } \overline{S})$$

sends the graded matrix factorization

$$\bigoplus_{i=1}^m R(d_i - h) \xrightarrow{\psi} \bigoplus_{i=1}^m R(e_i) \xrightarrow{\phi} \bigoplus_{i=1}^m R(d_i) \quad (3.1)$$

of f over R to the graded matrix factorization

$$\begin{array}{ccc} \bigoplus_{i=1}^m S(d_i - h) & \xrightarrow{\tilde{\psi} = \begin{pmatrix} \psi & -w \\ g & \phi \end{pmatrix}} & \bigoplus_{i=1}^m S(e_i) & \xrightarrow{\tilde{\phi} = \begin{pmatrix} \phi & w \\ -g & \psi \end{pmatrix}} & \bigoplus_{i=1}^m S(d_i) \\ \oplus & & \oplus & & \oplus \\ \bigoplus_{i=1}^m S(e_i - a) & & \bigoplus_{i=1}^m S(d_i - a) & & \bigoplus_{i=1}^m S(e_i + h - a), \end{array} \quad (3.2)$$

of F over S , where $\phi, \psi \in R$ are considered as elements of S by the injection $R \hookrightarrow S$.

Proof. Recall from [Eis80] that the matrix factorization (3.1) corresponds to the $\overline{R}$ -module $\overline{M} = \text{coker } \overline{\phi}$ through the 2-periodic projective resolution

$$\begin{array}{ccccccc} \cdots & \xrightarrow{\overline{\psi}} & \bigoplus_{i=1}^m \overline{R}(e_i - h) & \xrightarrow{\overline{\phi}} & \bigoplus_{i=1}^m \overline{R}(d_i - h) & \xrightarrow{\overline{\psi}} & \bigoplus_{i=1}^m \overline{R}(e_i) \\ & & & \searrow \overline{\phi} & \bigoplus_{i=1}^m \overline{R}(d_i) & \longrightarrow & \overline{M} \longrightarrow 0, \end{array} \quad (3.3)$$

where $\overline{\psi}$ and $\overline{\phi}$ are morphisms of $\overline{R}$ -modules induced by ψ and ϕ . By replacing each free $\overline{R}$ -modules in (3.3) with their $\overline{S}$ -free resolutions

$$0 \rightarrow \overline{S}(\ell - a) \xrightarrow{w} \overline{S}(\ell) \rightarrow \overline{R}(\ell) \rightarrow 0,$$

one obtains a projective resolution

$$\begin{array}{ccc} \cdots & \xrightarrow{\tilde{\psi}} & \begin{array}{c} \bigoplus_{i=1}^m \overline{S}(e_i - h) \\ \oplus \\ \bigoplus_{i=1}^m \overline{S}(d_i - a - h), \end{array} & \xrightarrow{\tilde{\phi}} & \begin{array}{c} \bigoplus_{i=1}^m \overline{S}(d_i - h) \\ \oplus \\ \bigoplus_{i=1}^m \overline{S}(e_i - a) \end{array} \\ & & \begin{array}{c} \bigoplus_{i=1}^m \overline{S}(e_i) \\ \oplus \\ \bigoplus_{i=1}^m \overline{S}(d_i - a) \end{array} & \xrightarrow{(\tilde{\phi} \quad w)} & \bigoplus_{i=1}^m \overline{S}(d_i) \longrightarrow \overline{M} \longrightarrow 0 \end{array}$$

of $\overline{M}$ as an $\overline{S}$ -module, where $\tilde{\phi}$ and $\tilde{\psi}$ are morphisms of $\overline{S}$ -modules induced by $\tilde{\phi}$ and $\tilde{\psi}$. This complex is 2-periodic except for the first two terms, and clearly corresponds to the matrix factorization (3.2), so that Proposition 3.1 is proved. $\square$

4 Morphisms between push-forwards

We keep the notations from Section 3. The following proposition gives Theorem 1.1 written in terms of matrix factorizations.

Proposition 4.1. *Let $\mathcal{E} = (\phi, \psi)$ and $\mathcal{E}' = (\phi', \psi')$ be matrix factorizations of f over R and $\mathcal{F} = \Phi_{\text{Hmf}}(\mathcal{E})$ and $\mathcal{F}' = \Phi_{\text{Hmf}}(\mathcal{E}')$ be their push-forwards. Then one has*

$$\text{Hom}(\mathcal{F}, \mathcal{F}') \cong \text{Hom}(\mathcal{E}, \mathcal{E}') \oplus \text{Hom}(\mathcal{E}', \mathcal{E}(r+h-a))^{\vee}[-d].$$

Proof. An element of $\text{Hom}(\mathcal{F}, \mathcal{F}')$ is represented by a pair (α, β) of morphisms

$$\alpha = \begin{pmatrix} \alpha_1 & \alpha_2 \\ \alpha_3 & \alpha_4 \end{pmatrix} : \begin{array}{c} \bigoplus_{i=1}^m S(d_i) \\ \oplus \\ \bigoplus_{i=1}^m S(e_i + h - a) \end{array} \longrightarrow \begin{array}{c} \bigoplus_{i=1}^m S(d'_i) \\ \oplus \\ \bigoplus_{i=1}^m S(e'_i + h - a) \end{array}$$

and

$$\beta = \begin{pmatrix} \beta_1 & \beta_2 \\ \beta_3 & \beta_4 \end{pmatrix} : \begin{array}{c} \bigoplus_{i=1}^m S(e_i) \\ \oplus \\ \bigoplus_{i=1}^m S(d_i - a) \end{array} \longrightarrow \begin{array}{c} \bigoplus_{i=1}^m S(e'_i) \\ \oplus \\ \bigoplus_{i=1}^m S(d'_i - a) \end{array}$$

making the diagram

$$\begin{array}{ccccc} \begin{array}{c} \bigoplus_{i=1}^m S(d_i - h) \\ \oplus \\ \bigoplus_{i=1}^m S(e_i - a) \end{array} & \xrightarrow{\tilde{\psi}} & \begin{array}{c} \bigoplus_{i=1}^m S(e_i) \\ \oplus \\ \bigoplus_{i=1}^m S(d_i - a) \end{array} & \xrightarrow{\tilde{\phi}} & \begin{array}{c} \bigoplus_{i=1}^m S(d_i) \\ \oplus \\ \bigoplus_{i=1}^m S(e_i + h - a) \end{array} \\ \alpha \downarrow & & \beta \downarrow & & \alpha \downarrow \\ \begin{array}{c} \bigoplus_{i=1}^m S(d'_i - h) \\ \oplus \\ \bigoplus_{i=1}^m S(e'_i - a) \end{array} & \xrightarrow{\tilde{\psi}'} & \begin{array}{c} \bigoplus_{i=1}^m S(e'_i) \\ \oplus \\ \bigoplus_{i=1}^m S(d'_i - a) \end{array} & \xrightarrow{\tilde{\phi}'} & \begin{array}{c} \bigoplus_{i=1}^m S(d'_i) \\ \oplus \\ \bigoplus_{i=1}^m S(e'_i + h - a) \end{array} \end{array}$$

commute; $\alpha \circ \tilde{\phi} = \tilde{\phi}' \circ \beta$ and $\beta \circ \tilde{\psi} = \tilde{\psi}' \circ \alpha$. The former condition can be written explicitly as the equality

$$\begin{pmatrix} \alpha_1 \phi - \alpha_2 g & \alpha_1 w + \alpha_2 \psi \\ \alpha_3 \phi - \alpha_4 g & \alpha_3 w + \alpha_4 \psi \end{pmatrix} = \begin{pmatrix} \phi' \beta_1 + w \beta_3 & \phi' \beta_2 + w \beta_4 \\ -g \beta_1 + \psi' \beta_3 & -g \beta_2 + \psi' \beta_4 \end{pmatrix} \quad (4.1)$$

where

$$\alpha \circ \tilde{\phi} = \begin{pmatrix} \alpha_1 & \alpha_2 \\ \alpha_3 & \alpha_4 \end{pmatrix} \begin{pmatrix} \phi & w \\ -g & \psi \end{pmatrix} = \begin{pmatrix} \alpha_1 \phi - \alpha_2 g & \alpha_1 w + \alpha_2 \psi \\ \alpha_3 \phi - \alpha_4 g & \alpha_3 w + \alpha_4 \psi \end{pmatrix}$$

and

$$\tilde{\phi}' \circ \beta = \begin{pmatrix} \phi' & w \\ -g & \psi' \end{pmatrix} \begin{pmatrix} \beta_1 & \beta_2 \\ \beta_3 & \beta_4 \end{pmatrix} = \begin{pmatrix} \phi' \beta_1 + w \beta_3 & \phi' \beta_2 + w \beta_4 \\ -g \beta_1 + \psi' \beta_3 & -g \beta_2 + \psi' \beta_4 \end{pmatrix},$$

and similarly as the equality

$$\begin{pmatrix} \beta_1 \psi + \beta_2 g & -\beta_1 w + \beta_2 \phi \\ \beta_3 \psi + \beta_4 g & -\beta_3 w + \beta_4 \phi \end{pmatrix} = \begin{pmatrix} \psi' \alpha_1 - w \alpha_3 & \psi' \alpha_2 - w \alpha_4 \\ g \alpha_1 + \phi' \alpha_3 & g \alpha_2 + \phi' \alpha_4 \end{pmatrix} \quad (4.2)$$

where

$$\beta \circ \tilde{\psi} = \begin{pmatrix} \beta_1 & \beta_2 \\ \beta_3 & \beta_4 \end{pmatrix} \begin{pmatrix} \psi & -w \\ g & \phi \end{pmatrix} = \begin{pmatrix} \beta_1\psi + \beta_2g & -\beta_1w + \beta_2\phi \\ \beta_3\psi + \beta_4g & -\beta_3w + \beta_4\phi \end{pmatrix}$$

and

$$\tilde{\psi}' \circ \alpha = \begin{pmatrix} \psi' & -w \\ g & \phi' \end{pmatrix} \begin{pmatrix} \alpha_1 & \alpha_2 \\ \alpha_3 & \alpha_4 \end{pmatrix} = \begin{pmatrix} \psi'\alpha_1 - w\alpha_3 & \psi'\alpha_2 - w\alpha_4 \\ g\alpha_1 + \phi'\alpha_3 & g\alpha_2 + \phi'\alpha_4 \end{pmatrix}$$

for the latter. Two morphisms (α, β) and (α', β') are homotopic if there exist morphisms

$$\xi = \begin{pmatrix} \xi_1 & \xi_2 \\ \xi_3 & \xi_4 \end{pmatrix} : \begin{array}{c} \bigoplus_{i=1}^m S(d_i) \\ \oplus \\ \bigoplus_{i=1}^m S(e_i + h - a) \end{array} \rightarrow \begin{array}{c} \bigoplus_{i=1}^m S(e'_i) \\ \oplus \\ \bigoplus_{i=1}^m S(d'_i - a) \end{array}$$

and

$$\eta = \begin{pmatrix} \eta_1 & \eta_2 \\ \eta_3 & \eta_4 \end{pmatrix} : \begin{array}{c} \bigoplus_{i=1}^m S(e_i) \\ \oplus \\ \bigoplus_{i=1}^m S(d_i - a) \end{array} \rightarrow \begin{array}{c} \bigoplus_{i=1}^m S(d'_i) \\ \oplus \\ \bigoplus_{i=1}^m S(e'_i + h - a) \end{array}$$

satisfying $\alpha - \alpha' = \tilde{\phi}' \circ \xi + \eta \circ \tilde{\psi}$ and $\beta - \beta' = \tilde{\psi}' \circ \eta + \xi \circ \tilde{\phi}$. The terms on the right hand sides can be explicitly written as

$$\begin{aligned} \tilde{\phi}' \circ \xi &= \begin{pmatrix} \phi' & w \\ -g & \psi' \end{pmatrix} \begin{pmatrix} \xi_1 & \xi_2 \\ \xi_3 & \xi_4 \end{pmatrix} = \begin{pmatrix} \phi'\xi_1 + w\xi_3 & \phi'\xi_2 + w\xi_4 \\ -g\xi_3 + \psi'\xi_3 & -g\xi_2 + \psi'\xi_4 \end{pmatrix}, \\ \eta \circ \tilde{\psi} &= \begin{pmatrix} \eta_1 & \eta_2 \\ \eta_3 & \eta_4 \end{pmatrix} \begin{pmatrix} \psi & -w \\ g & \phi \end{pmatrix} = \begin{pmatrix} \eta_1\psi + \eta_2g & -\eta_1w + \eta_2\phi \\ \eta_3\psi - \eta_4g & -\eta_3w + \eta_4\phi \end{pmatrix}, \\ \tilde{\psi}' \circ \eta &= \begin{pmatrix} \psi' & -w \\ g & \phi' \end{pmatrix} \begin{pmatrix} \eta_1 & \eta_2 \\ \eta_3 & \eta_4 \end{pmatrix} = \begin{pmatrix} \psi'\eta_1 - w\eta_3 & \psi'\eta_2 - w\eta_4 \\ g\eta_1 + \phi'\eta_3 & g\eta_2 + \phi'\eta_4 \end{pmatrix}, \\ \xi \circ \tilde{\phi} &= \begin{pmatrix} \xi_1 & \xi_2 \\ \xi_3 & \xi_4 \end{pmatrix} \begin{pmatrix} \phi & w \\ -g & \psi \end{pmatrix} = \begin{pmatrix} \xi_1\phi - \xi_2g & \xi_1w + \xi_2\psi \\ \xi_3\phi - \xi_4g & \xi_3w + \xi_4\psi \end{pmatrix}. \end{aligned}$$

Note that one can remove w -dependence of α_1 by choosing ξ_3 in a suitable way and achieve $\alpha_1 \in \text{Mat}_k(R)$. Similarly, w -dependence of α_2 , β_1 and β_2 can be removed by choosing ξ_4 , η_3 and η_4 respectively in a suitable way. Moreover, these operation of removing w -dependence can be performed independently, so that one can achieve $\alpha_1, \alpha_2, \beta_1, \beta_2 \in \text{Mat}_k(R)$ simultaneously.

Now the $(1, 1)$ -component of (4.1) gives the equation

$$\alpha_1\phi - \alpha_2g = \phi'\beta_1 + w\beta_3.$$

Since $\alpha_1, \beta_1, \phi, \phi' \in \text{Mat}_k(R)$ and $\alpha_2g, w\beta_3 \in w \text{Mat}_k(S)$ in the direct sum decomposition $\text{Mat}_k(S) = \text{Mat}_k(R) \oplus w \text{Mat}_k(S)$, one obtains

$$\alpha_1\phi = \phi'\beta_1 \quad \text{and} \quad \beta_3 = -\frac{g}{w}\alpha_2. \quad (4.3)$$

Similarly, from the $(1, 2)$ -component of (4.1), one obtains

$$\alpha_1 w + \alpha_2 \psi = \phi' \beta_2 + w \beta_4$$

which, together with $\alpha_2, \beta_2, \psi, \phi' \in \text{Mat}_k(R)$, implies

$$\alpha_2 \psi = \phi' \beta_2 \quad \text{and} \quad \beta_4 = \alpha_1. \quad (4.4)$$

The same argument for $(1, 1)$ - and $(1, 2)$ -components of (4.2) gives

$$\beta_1 \psi = \psi' \alpha_1 \quad \text{and} \quad \alpha_3 = -\frac{g}{w} \beta_2 \quad (4.5)$$

and

$$\beta_2 \phi = \psi' \alpha_2 \quad \text{and} \quad \alpha_4 = \beta_1 \quad (4.6)$$

respectively. This determines α and β completely from $(\alpha_1, \alpha_2, \beta_1, \beta_2)$ satisfying

$$\alpha_1 \phi = \phi' \beta_1, \quad \alpha_2 \psi = \phi' \beta_2, \quad \beta_1 \psi = \psi' \alpha_1, \quad \beta_2 \phi = \psi' \alpha_2. \quad (4.7)$$

Other components of (4.1) and (4.2) are automatically satisfied for α and β satisfying (4.3), (4.4), (4.5), and (4.6). This shows that any element of $\text{Hom}(\mathcal{F}, \mathcal{F}')$ can be represented by a pair (α, β) of morphisms coming from a representative (α_1, β_1) of $\text{Hom}(\mathcal{E}, \mathcal{E}')$ and a representative (α_2, β_2) of $\text{Hom}(\mathcal{E}[1](-a), \mathcal{E}')$.

If two morphisms from $\mathcal{F}$ to $\mathcal{F}'$ represented by pairs (α, β) and (α', β') coming from representatives (α_1, β_1) and (α'_1, β'_1) of $\text{Hom}(\mathcal{E}, \mathcal{E}')$ and representatives (α_2, β_2) and (α'_2, β'_2) of $\text{Hom}(\mathcal{E}[1](-a), \mathcal{E}')$ are homotopic, then one has two morphisms ξ and η satisfying $\alpha - \alpha' = \tilde{\phi}' \circ \xi + \eta \circ \tilde{\psi}$ and $\beta - \beta' = \tilde{\psi}' \circ \eta + \xi \circ \tilde{\phi}$. By looking at the $\text{Mat}_{2k}(R)$ component of these equations with respect to the direct sum decomposition $\text{Mat}_{2k}(S) = \text{Mat}_{2k}(R) \oplus w \text{Mat}_{2k}(S)$, one finds that (α_1, β_1) and (α'_1, β'_1) are homotopic over R , and (α_2, β_2) and (α'_2, β'_2) are homotopic over R as well. It follows that one has an isomorphism

$$\text{Hom}(\mathcal{F}, \mathcal{F}') \cong \text{Hom}(\mathcal{E}, \mathcal{E}') \oplus \text{Hom}(\mathcal{E}[1](-a), \mathcal{E}')$$

of spaces of morphisms. Graded Auslander-Reiten duality [AR87] implies Serre duality

$$\text{Hom}(\mathcal{M}, \mathcal{N}) \cong \text{Hom}(\mathcal{N}, \mathcal{M}(r+h)[d-1])^\vee$$

in the graded stable derived category of $\overline{R}$ [IT, Corollary 2.5], so that

$$\text{Hom}(\mathcal{E}[1](-a), \mathcal{E}') \cong \text{Hom}(\mathcal{E}', \mathcal{E}(r+h-a))^\vee[-d]$$

and Proposition 4.1 is proved. $\square$

References

- [AR87] Maurice Auslander and Idun Reiten, *Almost split sequences for $\mathbf{Z}$ -graded rings*, Singularities, representation of algebras, and vector bundles (Lambrecht, 1985), Lecture Notes in Math., vol. 1273, Springer, Berlin, 1987, pp. 232–243. MR 915178 (89b:13031)

- [BH93] Per Berglund and Tristan Hübsch, *A generalized construction of mirror manifolds*, Nuclear Phys. B **393** (1993), no. 1-2, 377–391. MR MR1214325 (94k:14031)
- [Bor] Lev A. Borisov, *Berglund-Hübsch mirror symmetry via vertex algebras*, arXiv:1007.2633.
- [Buc87] Ragnar-Olaf Buchweitz, *Maximal Cohen-Macaulay modules and Tate-cohomology over Gorenstein rings*, Available from <https://tspace.library.utoronto.ca/handle/1807/16682>, 1987.
- [CR11] Alessandro Chiodo and Yongbin Ruan, *LG/CY correspondence: the state space isomorphism*, Adv. Math. **227** (2011), no. 6, 2157–2188. MR 2807086
- [Eis80] David Eisenbud, *Homological algebra on a complete intersection, with an application to group representations*, Trans. Amer. Math. Soc. **260** (1980), no. 1, 35–64. MR MR570778 (82d:13013)
- [FU] Masahiro Futaki and Kazushi Ueda, *Homological mirror symmetry for singularities of type D*, arXiv:1004.0078, to appear in Mathematische Zeitschrift.
- [FU11] ———, *Homological mirror symmetry for Brieskorn-Pham singularities*, Selecta Math. (N.S.) **17** (2011), no. 2, 435–452.
- [Hap91] Dieter Happel, *On Gorenstein algebras*, Representation theory of finite groups and finite-dimensional algebras (Bielefeld, 1991), Progr. Math., vol. 95, Birkhäuser, Basel, 1991, pp. 389–404. MR 1112170 (92k:16022)
- [IT] Osamu Iyama and Ryo Takahashi, *Tilting and cluster tilting for quotient singularities*, arXiv:1012.5954.
- [KMU12] Masanori Kobayashi, Makiko Mase, and Kazushi Ueda, *A note on exceptional unimodal singularities and K3 surfaces*, Internat. Math. Res. Notices (2012), no. rns098, 1–26.
- [Kon95] Maxim Kontsevich, *Homological algebra of mirror symmetry*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994) (Basel), Birkhäuser, 1995, pp. 120–139. MR MR1403918 (97f:32040)
- [Kra] Marc Krawitz, *FJRW rings and Landau-Ginzburg mirror symmetry*, arXiv:0906.0796.
- [Kra05] Henning Krause, *The stable derived category of a Noetherian scheme*, Compos. Math. **141** (2005), no. 5, 1128–1162. MR MR2157133 (2006e:18019)
- [NU] Yuichi Nohara and Kazushi Ueda, *Homological mirror symmetry for the quintic 3-fold*, arXiv:1103.4956, to appear in Geometry and Topology.
- [Orl04] D. O. Orlov, *Triangulated categories of singularities and D-branes in Landau-Ginzburg models*, Tr. Mat. Inst. Steklova **246** (2004), no. Algebr. Geom. Metody, Svyazi i Prilozh., 240–262. MR MR2101296

- [Orl09] Dmitri Orlov, *Derived categories of coherent sheaves and triangulated categories of singularities*, Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II, Progr. Math., vol. 270, Birkhäuser Boston Inc., Boston, MA, 2009, pp. 503–531. MR 2641200 (2011c:14050)
- [Sei02] Paul Seidel, *Fukaya categories and deformations*, Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002) (Beijing), Higher Ed. Press, 2002, pp. 351–360. MR MR1957046 (2004a:53110)
- [Sei08] ———, *Fukaya categories and Picard-Lefschetz theory*, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), Zürich, 2008. MR MR2441780
- [Sei11] ———, *Homological mirror symmetry for the quartic surface*, math.AG/0310414, 2011.
- [She11] Nick Sheridan, *Homological mirror symmetry for Calabi-Yau hypersurfaces in projective space*, arXiv:1111.0632, 2011.
- [Tak10] Atsushi Takahashi, *Weighted projective lines associated to regular systems of weights of dual type*, New developments in algebraic geometry, integrable systems and mirror symmetry (RIMS, Kyoto, 2008), Adv. Stud. Pure Math., vol. 59, Math. Soc. Japan, Tokyo, 2010, pp. 371–388. MR 2683215

Kazushi Ueda

Department of Mathematics, Graduate School of Science, Osaka University, Machikaneyama 1-1, Toyonaka, Osaka, 560-0043, Japan.

e-mail address : kazushi@math.sci.osaka-u.ac.jp